arXiv:hep-ph/0112293v1 21 Dec 2001

BUTP-2001/23
BUHE-2001-07

HadAtom01

Workshop on Hadronic Atoms
Institut für Theoretische Physik, Universität Bern
Sidlerstrasse 5, CH-3012 Bern, Schweiz
October 11 - 12, 2001

Jürg Gasser[1]
Institut für Theoretische Physik, Universität Bern, CH-3012 Bern, Schweiz

Akaki Rusetsky[2]
Institut für Theoretische Physik, Universität Bern, CH-3012 Bern, Schweiz
HEPI, Tbilisi State University, 380036 Tbilisi, Georgia

Jürg Schacher[3]
Laboratorium für Hochenergiephysik, Universität Bern, CH-3012 Bern, Schweiz

Abstract

These are the proceedings of the workshop "HadAtom01", held at the Institut für Theoretische Physik, Universität Bern, October 11 - 12, 2001. The main topics discussed at the workshop were the physics of hadronic atoms and in this context recent results in experiment and theory. Included here are the list of participants, the scientific program and a short contribution from each speaker.

---

[1] E-mail: gasser@itp.unibe.ch
[2] E-mail: rusetsky@itp.unibe.ch
[3] E-mail: schacher@lhep.unibe.ch

1 Introduction

The workshop "HadAtom01" took place at the Institut für Theoretische Physik, Universität Bern on October 11-12, 2001. It was the third in a series of workshops on bound states, in particular hadronic atoms: the previous ones were held in Dubna, Russia (May 1998) [1], and in Bern (October 1999) [2]. The meeting was attended by about 50 physicists, and results were presented by 25 participants.

The topics of the workshop included

- Hadronic atoms, in particular their
  - Production
  - Interaction with matter
  - Energy levels
  - Decays
- Meson-meson and meson-baryon scattering
- Experiments
  - DIRAC at CERN
  - DEAR at DAFNE
  - PSI (Pionic Hydrogen Collaboration)
  - Others
- $K_{\ell 4}$ decays

The talks were devoted to recent experimental and theoretical progress in the investigations of hadronic atoms. Among the highlights of the workshop were the presentation of preliminary results of the DIRAC collaboration on the measurement of the lifetime of pionium, as well as the first measurement of kaonic nitrogen by the DEAR collaboration.

The speakers have provided a one-page abstract of the presentations as well as a list of the most relevant references - these contributions are collected below. We also display a list of the participants with their e-mail, and the scientific program of the workshop.

Acknowledgement

We would like to thank all participants for their effort to travel to Bern and for making "HadAtmo01" an exciting and lively meeting. We furthermore thank our secretaries, Ruth Bestgen and Ottilia Hänni, for their excellent support, Manuel Walker for designing the poster of the workshop, and Christoph Haefeli, Andrea Macrina, Bernhard Scheuner, Julia Schweizer and Peter Zemp for their help during the workshop. Last but not least, we thank our colleagues from the organizing committee (Valery Lyubovitskij, Leonid Nemenov, Hagop Sazdjian, and Dirk Trautmann) for their invaluable contribution in structuring the meeting. This work was supported in part by the Swiss National Science Foundation, and by TMR, BBW-Contract No. 97.0131 and EC-Contract No. ERBFMRX-CT980169 (EURODAΦNE).

Bern, December 2001

Jürg Gasser, Akaki Rusetsky and Jürg Schacher

---

[1] Proceedings of the International Workshop "Hadronic Atoms and Positronium in the Standard Model", Dubna, 26-31 May 1998, ed. M.A. Ivanov, A.B. Arbuzov, E.A. Kuraev, V.E. Lyubovitskij, A.G. Rusetsky.

[2] J. Gasser, A. Rusetsky and J. Schacher, arXiv:hep-ph/9911339.

2 List of participants

1.	Afanasev Leonid (Dubna)	leonid.afanasev@cern.ch
2.	Aste Andreas (Basel)	andreas.aste@unibas.ch
3.	Brekhovskikh Valeri (Protvino)	valeri.brekhovskikh@cern.ch
4.	Büttiker Paul (Jülich)	p.buettiker@fz-juelich.de
5.	Colangelo Gilberto (Zürich)	gilberto@physik.unizh.ch
6.	Cribeiro Alberto (Santiago)	alberto.cribeiro@rai.usc.es
7.	Cuplov Vesna (Marseille)	cuplov@cpt.univ-mrs.fr
8.	Curceanu [Petrascu] Catalina (Frascati)	catalina@lnf.infn.it
9.	Drijard Daniel (CERN)	daniel.drijard@cern.ch
10.	Egger Jean-Pierre (Neuchâtel)	jean-pierre.egger@unine.ch
11.	Eiras Dolors (Barcelona)	dolors@ecm.ub.es
12.	Gasser Juerg (Bern)	gasser@itp.unibe.ch
13.	Gianotti Paola (Frascati)	paola.gianotti@lnf.infn.it
14.	Goldin Daniel (Basel)	daniel.goldin@cern.ch
15.	Gotta Detlev (Jülich)	d.gotta@fz-juelich.de
16.	Guaraldo Carlo (Frascati)	guaraldo@lnf.infn.it
17.	Hasenfratz Peter (Bern)	hasenfra@itp.unibe.ch
18.	Heim Thomas (Basel)	thomas.heim@unibas.ch
19.	Hencken Kai (Basel)	k.hencken@unibas.ch
20.	Hennebach Maik (Köln)	m.hennebach@fz-juelich.de
21.	Jensen Thomas (PSI)	thomas.jensen@psi.ch
22.	Knecht Marc (Marseille)	knecht@cpt.univ-mrs.fr
23.	Kubis Bastian (Jülich)	b.kubis@fz-juelich.de
24.	Lanaro Armando (CERN)	armando.lanaro@cern.ch
25.	Lipartia Edisher (Lund, Dubna & Tbilisi)	lipartia@thep.lu.se
26.	Ludhova Livia (Fribourg)	livia.ludhova@unifr.ch
27.	Lyubovitskij Valery (Tübingen & Tomsk)	valeri.lyubovitskij@uni-tuebingen.de
28.	Meißner Ulf-G. (Jülich)	u.meissner@fz-juelich.de
29.	Nehme Abbass (Marseille)	nehme@cpt.univ-mrs.fr
30.	Nemenov Leonid (CERN & Dubna)	leonid.nemenov@cern.ch
31.	Nyffeler Andreas (Marseille)	nyffeler@cpt.univ-mrs.fr
32.	Pancheri Giulia (Frascati)	pancheri@lnf.infn.it
33.	Pentia Mircea (Bucharest)	pentia@ifin.nipne.ro
34.	Rasche Günther (Zürich)	rasche@physik.unizh.ch
35.	Rusetsky Akaki (Bern & Tbilisi)	rusetsky@itp.unibe.ch
36.	Sainio Mikko (Helsinki)	mikko.sainio@helsinki.fi
37.	Santamarina Cibrán (Santiago)	cibran.santamarina.rios@cern.ch
38.	Sazdjian Hagop (Orsay)	sazdjian@ipno.in2p3.fr
39.	Schacher Juerg (Bern)	schacher@lhep.unibe.ch
40.	Schumann Marc (Basel)	marc.schumann@unibas.ch
41.	Schütz Christian (Basel)	christian.schuetz@cern.ch
42.	Schweizer Julia (Bern)	schweize@itp.unibe.ch
43.	Simons Leopold (PSI)	leopold.simons@psi.ch
44.	Tarasov Alexander (Dubna)	avt@mpimail.mpi-hd.mpg.de
45.	Tauscher Ludwig (Basel)	ludwig.tauscher@cern.ch
46.	Trautmann Dirk (Basel)	dirk.trautmann@unibas.ch
47.	Voskresenskaya Olga (Dubna)	voskr@mpimail.mpi-hd.mpg.de
48.	Zemp Peter (Bern)	zemp@itp.unibe.ch
49.	Zmeskal Johann (Vienna)	zmeskal@amuon.imep.univie.ac.at

3 Scientific program

Page

C. Santamarina
DIRAC (PS-212) experiment at CERN 6

L. Afanasyev, A. Tarasov and O.Voskresenskaya
Spectrum of pions from $\pi^+\pi^-$-atom breakup (Born and Glauber approximations) 7

L.L. Nemenov
Possibility to measure the energy splitting between $2S$ and $2P$ states in $\pi^+\pi^-$ atoms 8

J. Gasser, I. Mgeladze, **A. Rusetsky** and I. Scimemi
Hadronic atoms: review of theoretical aspects 9

V.E. Lyubovitskij
Spectrum and decays of $\pi^+\pi^-$ and π^-p atoms 10

G. Rasche
Electromagnetic corrections to the lifetime of pionium 11

E. Lipartia, V. Lyubovitskij and A. Rusetsky
Hadronic potentials from effective field theory 12

T.A. Heim, K. Hencken, M. Schumann, D. Trautmann and G. Baur
Distribution of pions from breakup of pionium 13

M. Schumann, T. Heim, K. Hencken, D. Trautmann and G. Baur
Higher order corrections to bound-bound excitation cross sections of pionium 14

T. Heim, **K. Hencken**, M. Schumann, D. Trautmann and G. Baur
Calculation of electromagnetic breakup of pionium 15

A. Lanaro
πK atom: Observation and lifetime measurement with DIRAC 16

H. Sazdjian
Can one extract $m_s/\hat{m}$ from πK atom properties? 17

B. Ananthanarayan, **P. Büttiker** and B. Moussallam
πK scattering and dispersion relations 18

B. Kubis and U.-G. Meißner
Isospin violation in pion-kaon scattering 19

C. Curceanu (Petrascu)
The first measurement of kaonic nitrogen with DEAR at DAFNE 20

P. Hasenfratz
The quark condensate on the lattice 21

G. Colangelo
Chiral and lattice calculations of the $\pi\pi$ scattering lengths 22

G. Colangelo, **J. Gasser** and H. Leutwyler
The quark condensate from K_{e4} decays 23

M. Knecht
Radiative corrections to semileptonic decays 24

V. Cuplov
Isospin violation in $K_{\ell 4}$ decays 25

M. Knecht and **A. Nyffeler**
Resonance estimates of low-energy constants in chiral Lagrangians and QCD short-distance constraints 26

M. Hennebach
The new pionic hydrogen experiment at PSI 27

T. Jensen and V.E. Markushin
Atomic cascade in hadronic atoms 28

M.E. Sainio
Pion-nucleon analysis 29

N. Fettes and **U.-G. Meißner**
Theory of low-energy pion-nucleon scattering 30

DIRAC (PS-212) experiment at CERN

C. Santamarina[†] [4] **(on behalf of DIRAC collaboration)**

[†] *Departamento de Física de Partículas, Universidade de Santiago de Compostela, Campus Universitario Sur, 15782- Santiago de Compostela, Spain.*

The theoretical results in the low energy regime of hadronic physics have reached an accuracy level which should be tested with new and precise experimental measurements. In particular, a prediction of $2.9 \pm 0.1 \times 10^{-15}$ seconds [1] has been made for the lifetime of the ground state of pionium, the $\pi^+\pi^-$ electromagnetic atom, within the framework of Chiral Perturbation Theory.

DIRAC-PS 212 [2] is a fixed target experiment, currently running at CERN PS, which aims to measure the pionium lifetime with a 10% accuracy analyzing the low relative momentum region of $\pi^+\pi^-$ pairs spectrum ($Q < 25 MeV/c$) resulting from proton target collisions [3].

The DIRAC spectrometer is a double arm telescope with a central magnet. It is optimized to detect events with two oppositely charged particles with low relative momentum. The detectors provide an efficient particle identification to exclude events with proton, electron or muon contamination.

The evidence of pairs from atomic break-up was first found in Summer 2000. The analysis of 2000 and 2001 data has given us around 5600 atoms mainly with a Titanium (1800 atoms) 175 μm and a Nickel (3600 atoms) 94 μm targets.

With the analyzed data a preliminary result, considering statistical errors only, has been obtained both with Nickel and Titanium data. The combined analysis of both samples gives us a lifetime of $3.6^{+0.9}_{-0.7} \times 10^{-15} s$, however a detailed analysis of systematic errors is yet to be performed.

Sample Target/Year	Lifetime ($\times 10^{-15} s$)	Statistical error
Ni 2000	$2.8^{+1.1}_{-0.8}$	±34%
Ti 2000/1	$5.4^{+1.5}_{-1.3}$	±26%
Ti+Ni	$3.6^{+0.9}_{-0.7}$	±22%

Table 1: Preliminary results with 2000 and 2001 data. Only statistical errors have been analyzed.

[1] J. Gasser, V.E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008.

[2] B. Adeva *et al.*, CERN-SPSLC-95-1 ; SPSLC-P-284 (1994).

[3] L.L. Nemenov, Yad. Fiz. **41** (1985) 980; Sov. J. Nucl. Phys. **41** (1985) 629.

---

[4] Now at Basel University.

Spectrum of pions from $\pi^+\pi^-$-atom breakup (Born and Glauber approximations)

L. Afanasyev[1], **A. Tarasov**[1,2] **and O.Voskresenskaya**[1]

[1] *Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia*
[2] *Institute for Theoretical Phisics, University of Heidelberg, Philosophenweg 19, D-69120, Heidelberg, Germany*

Momentum and angular distributions in $\pi^+\pi^-$-pairs from the $\pi^+\pi^-$-atom ($A_{2\pi}$) breakup at the electromagnetic interactions with target atoms have been considered in the Born and Glauber approximations. Exact analytical expressions for the spectra for the initial states with orbital quantum number $l = 0$ (nS states) have obtained in the first Born approximation.



n=1 n=10

In the figure the distributions over the relative momentum of pions p and angle θ with respect to the total momentum of atom are shown for the initial states with $n = 1$ and $n = 10$ for the nickel target.

The formulas for the discussed spectra for the initial ground states of $A_{2\pi}$ which takes into account muli-photon exchange processes have been obtained in the following approach. It is shown that due to the small size of $A_{2\pi}$ and its big mass the interaction with an unscreened target nuclear dominantly contributes to the multi-photon exchange processes. The amplitude of the atom interaction with unscreened Coulomb potential can be expressed in a closed analytical form in the Glauber approximation. Thus one can consider the target atom as the screened Coulomb potential for accounting the single-photon exchange processes in the Born approximation and as unscreened one for multi-photon processes.

Possibility to measure the energy splitting between 2S and 2P states in $\pi^+\pi^-$ atoms

L.L. Nemenov

CERN, CH-1211, Geneva 23, Switzerland and Dubna (JINR), Russia

The difference ΔE_n between the energy of atomic nS and nP states of $A_{2\pi}$ includes the contributions of the vacuum polarisation ΔE_n^{vac}, and of the strong interaction ΔE_n^s effects, where $\Delta E_n^s \sim 2a_0 + a_2$, and a_0 and a_2 are the S-wave $\pi\pi$ scattering lengths with isotopic-spin quantum numbers 0,2 [1,2,3].

The value of ΔE_n^{vac} is well known from QED calculations [1,2,4]. For this reason the measurement of ΔE_n will give, in a model independent way, the value of $2a_0 + a_2$ [5,6].

The method for measuring ΔE_n is presented [7,8]. The same method allows also to measure the energy splitting (ΔE_n) for atoms consisting of π and K mesons, and to obtain the value of $2a_{1/2} + a_{3/2}$, where $a_{1/2}$ and $a_{3/2}$ are the S-wave πK scattering lengths with isotopic-spin 1/2 and 3/2, respectively.

[1] A. Gashi et al., Nucl. Phys. A **628** (1999) 101.
[2] A. Rusetsky, private communication.
[3] G. Efimov et al., Sov. J. Nucl. Phys. **44** (1986) 296.
[4] A. Karimkhodzhaev and R.N. Faustov, Sov. J. Nucl. Phys. **29** (1979) 232.
[5] G. Colangelo, J. Gasser and H. Leutwyler, Phys. Lett. B **488** (2000) 261.
[6] M. Knecht et al., Nucl. Phys. B **457** (1995) 513.
[7] L.L. Nemenov, Sov. J. Nucl. Phys.**41** (1985) 629.
[8] L.L. Nemenov and V.D. Ovsiannikov, Phys. Lett. B **514** (2001) 247.

Hadronic atoms: review of theoretical aspects

J. Gasser, I. Mgeladze, A. Rusetsky and I. Scimemi

Institute for Theoretical Physics, University of Bern, Sidlerstrasse 5, CH-3012, Bern, Switzerland

In order to fully exploit the high-precision experimental data on hadronic atoms which is available at present and which will be provided in the future, it is imperative to design a theoretical framework that describes this type of bound systems to the accuracy that matches the experimental precision. The Deser-type relations which are then used to extract the "purely strong" scattering lengths from the measured values of the strong energy shift in the ground state, and the partial decay width into hadronic channels, contain isospin-breaking corrections which depend on the fine structure constant α and on the quark mass difference $m_d - m_u$. The following questions arise.

i) The scattering lengths are calculated in QCD at $m_u = m_d$. How is this theory related to a framework that includes isospin violating effects, $SU(3)_c \times U(1)_{em}$? [Note that, in view of the announced accuracy 0.3 % in the future measurement of the energy shift of the $\pi^- p$ atom by Pionic Hydrogen collaboration at PSI, the question of the precise definition of the pure QCD limit is not an academic one.]

ii) With a given definition of the pure QCD limit, how does one calculate the isospin-breaking corrections to the bound-state observables?

The answers are as follows.

ad i) In two-flavour QCD, the quark mass $m_u = m_d$ and the strong coupling constant may be fixed from the requirement that the pion and nucleon masses are equal - by convention - to the experimental values of the charged pion and the proton masses, respectively. In the effective theory of $SU(3)_c \times U(1)_{em}$, the values of the strong LECs stay put [1].

ad ii) We have developed a general theory of hadronic atoms, based on a merger of non-relativistic effective Lagrangian techniques and ChPT [3-6]. This theory has been successfully applied to the calculation of the isospin-breaking corrections in the $\pi^+\pi^-$ atom decay width [3,4,6], and the energy shift of the $\pi^- p$ atom [5]. For related work, see [7]. Applications to other systems are in preparation [8].

[1] J. Gasser, I. Mgeladze, A. Rusetsky and I. Scimemi, in preparation. This reference reviews earlier work on the subject. We have profited from unpublished notes by H. Leutwyler, whom we thank in addition for informative discussions on the subject.

[3] A. Gall, J. Gasser, V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **462** (1999) 335.

[4] J. Gasser, V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **471** (1999) 244 (1999).

[5] V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **494** (2000) 9.

[6] J. Gasser, V.E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008.

[7] X. Kong and F. Ravndal, Phys. Rev. D 61 (2000) 077506 ; D. Eiras and J. Soto, Phys. Lett. B 491 (2000) 101, and references cited therein.

[8] M. Schmid, J. Schweizer and P. Zemp, work in progress.

Spectrum and decays of $\pi^+\pi^-$ and $\pi^- p$ atoms [1]

V. E. Lyubovitskij
Institut für Theoretische Physik, Universität Tübingen,
Auf der Morgenstelle 14, D-72076 Tübingen, Germany

In order to carry out the precision experimental tests of QCD we proposed and developed *effective theory* [2,3] for the description of measured characteristics of hadronic atoms. We have derived the expressions for the decay width $\Gamma_{2\pi^0}$ of the $\pi^+\pi^-$ atom [2] and strong energy shift of the $\pi^- p$ atom [3] in the framework of QCD (including photons) by use of effective field theory techniques. The results obtained contains all terms at leading and next-to-leading order in the isospin breaking parameters $\alpha \simeq 1/137$ and $m_u - m_d$. On the basis of these formulae, a numerical analysis was carried out in Refs. [2,3] at order $e^2 p^2$ in chiral perturbation theory (ChPT).

The $\pi^+\pi^-$ atom decay problem is completely solved [2]. Numerical analysis in ChPT based on recent update results for the S-wave $\pi\pi$ scattering lengths a_0 and a_2 gives

$$\tau \doteq \Gamma_{2\pi^0}^{-1} = (2.9 \pm 0.1) \times 10^{-15}\ \mathrm{s}\,. \tag{1}$$

Numerical analysis of the $\pi^- p$ atom strong energy shift is in progress because the effective electromagnetic coupling constant f_1 is still unknown.

The perturbative chiral quark model [4] was applied to estimate the electromagnetic $O(p^2)$ low-energy constants of the ChPT effective Lagrangian that define the electromagnetic mass shifts of nucleons and first-order (e^2) radiative corrections to the πN scattering amplitude [5]. It was found that the value of f_2 constant is consistent with model-independent result obtained by Gasser and Leutwyler [6]. Using obtained value for constant f_1 the leading isospin-breaking correction to the strong energy shift of the $\pi^- p$ atom in the $1s$ state was calculated. The result $\delta_\epsilon = -2.8 \times 10^{-2}$ is comparable to a prediction based on a potential model for the πN scattering [7]: $\delta_\epsilon = -2.1 \times 10^{-2}$.

Acknowledgements. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, grant FA67/25-1).

[1] Work done in collaboration with J. Gasser, A. Rusetsky, A. Gall [2,3] and Th. Gutsche, A. Faessler, R. Vinh Mau [4,5].

[2] A. Gall, J. Gasser, V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **462** (1999) 335; J. Gasser, V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **471** (1999) 244; J. Gasser, V.E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008.

[3] V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **494** (2000) 9.

[4] V.E. Lyubovitskij, T. Gutsche, A. Faessler and E.G. Drukarev, Phys. Rev. D **63** (2001) 054026.

[5] V.E. Lyubovitskij, T. Gutsche, A. Faessler and R. Vinh Mau, Phys. Lett. B **520** (2001) 204.

[6] J. Gasser and H. Leutwyler, Phys. Rep. **87** (1982) 77.

[7] D. Sigg, et al, Nucl. Phys. A **609** (1996) 310.

Electromagnetic corrections to the lifetime of pionium

G. Rasche

Institute for Theoretical Physics, University of Zürich, Winterthurerstr. 190, CH-8057, Zürich, Switzerland

We have made use of the Deser formula for the two-channel system as derived in [1]. Including a tiny correction for vacuum polarisation etc. in the numerical constant, we get from [2]:

$$\tau\,(\mathrm{fs}) = \frac{0.09474}{(a_{0\mathrm{c}}\,(\mathrm{fm}))^2}\ . \tag{1}$$

Here a_{0c} is the value of the $\pi^+\pi^- \to \pi^0\pi^0$ transition element of the **K**-matrix of scattering theory at the $\pi^+\pi^-$ threshold. It consists of a hadronic part $a^h_{0c}(\mu_0)$ and an electromagnetic correction $\Delta a_{0c}(\mu_0)$:

$$a_{0c} = a^h_{0c}(\mu_0) + \Delta a_{0c}(\mu_0)\ . \tag{2}$$

Since the pion mass for the hadronic limit is very close to the neutral pion mass μ_0 it follows that the hadronic starting point for the true electromagnetic corrections is μ_0. We have indicated this explicitly on the lhs of (2).

The value for $a^h_{0c}(\mu_0)$ is taken from [3]; they use $a^0 - a^2 = 0.265(4)$. Taking into account that this value corresponds to a hadronic starting point with pion mass μ_c and is given in units of μ_c^{-1} one gets (see [2])

$$a^h_{0c}(\mu_0) = \frac{\sqrt{2}}{3}(a^2(\mu_0) - a^0(\mu_0)) = -0.170(3)\,\mathrm{fm}\ . \tag{3}$$

The electromagnetic correction $\Delta a_{0c}(\mu_0)$ is calculated using a relativised two-channel Schrödinger equation with two isospin invariant energy independent potentials $V^I(r), I = 0, 2$. They are constructed by fitting an analytical ansatz to the two-loop hadronic ChPT scattering phase $\delta^I(q), I = 0, 2$ of [4], for both values μ_0 and μ_c of the pion mass. Solving the coupled Schrödinger equation including the Coulomb interaction and the mass difference $\mu_c - \mu_0$, as well as taking into account the results of [5], we arrive at

$$\Delta a_{0c}(\mu_0) = 0.010(1)\,\mathrm{fm}\ . \tag{4}$$

Inserting these results into (1) we get

$$\tau = 2.92(15)\,\mathrm{fs}\ . \tag{5}$$

This agrees with [3].

[1] G. Rasche and W.S. Woolcock, Nucl.Phys. A **381** (1982) 405.

[2] A. Gashi, G.C. Oades, G. Rasche and W.S.Woolcock, arXiv:hep-ph/0108116, to appear in Nucl. Phys. A.

[3] J. Gasser, V.E. Lyubovitskij, A. Rusetsky and A. Gall, Phys. Rev. D **64** (2001) 016008.

[4] J. Bijnens, G. Colangelo, G. Ecker, J. Gasser and M.E. Saino, Phys.Lett. B **374** (1996) 210; Nucl. Phys. B **508** (1997) 263 and erratum ibid. B **517** (1998) 693; G. Colangelo, private communication.

[5] M. Knecht and R.Urech, Nucl. Phys. B **519** (1998) 329.

Hadronic potentials from effective field theory

E. Lipartia[1,2,3], **V. Lyubovitskij**[4,5] **and A. Rusetsky**[3,6]

[1] *Department of Theoretical Physics 2, Lund University, Solvegatan 14A, S22362 Lund, Sweden*
[2] *LIT, Joint Institute for Nuclear Research, 141980, Dubna, Russia*
[3] *HEPI, Tbilisi State University, University st. 9, 380086 Tbilisi, Georgia*
[4] *Institute of Theoretical Physics, University of Tübingen, Auf der Morgenstelle 14, D-72076, Tübingen, Germany*
[5] *Department of Physics, Tomsk State University, 634050 Tomsk, Russia*
[6] *Institute for Theoretical Physics, University of Bern, Sidlerstrasse 5, CH-3012, Bern, Switzerland*

The problem of the evaluation of isospin symmetry breaking effects to the hadronic atom observables has been addressed in the framework of field-theoretical approaches, as well as in the potential models. At the same time, the results obtained within the potential model do not always agree with the results of field-theoretical approaches. The main reason of discrepancy between the predictions of both approaches is that, in general, the potential model does not take into account the full content of isospin-breaking effects in ChPT. Even if these predictions agree in a particular case (e.g., like the latest prediction of the potential model for the lifetime of $\pi^+\pi^-$ atom [1]), without an explicit matching to ChPT in the isospin-breaking phase the treatment of the problem in the potential model can not be regarded as systematic. In the view of the fact that the potential approach is widely used to evaluate the isospin-breaking effects in scattering process - to analyze the results of experimental measurements - it is very important to construct the potential which includes the full content of isospin-breaking effects.

The constructive algorithm for the derivation of the hadronic potential from ChPT is considered in detail in our paper [2]. Here we provide only a brief outline of the procedure. The derivation of the isospin-breaking part of the short-range strong potential is based on the universality conjecture, which states that the relation between the bound-state characteristics and scattering amplitudes is the same in the field theory and in the potential model. The universality conjecture provides us with the matching condition for the isospin-breaking part of hadronic potential, assuming the isospin-symmetric part of the potential is given. Further, the matching condition imposes rather loose constraints on the potential. This looseness reflects the freedom of the choice of regularization of pointlike singularities which emerges if the potential is derived from a local field theory. For this reason, the shape of the potential does not bear a physical information, and using this freedom, one may absorb all information about the isospin-breaking effects in the couplings of the short-range potential which, in its turn, are determined from the matching condition [2].

Despite the unrestricted freedom in the choice of the shape of the short-range part of the potential, the full potential that we construct, reproduces, by construction, the threshold amplitude and the bound-state characteristics of hadronic atoms at the first non-leading order in isospin-breaking.

[1] G. Rasche, these proceedings.

[2] E. Lipartia, V.E. Lyubovitskij and A. Rusetsky, arXiv:hep-ph/0110186.

Distribution of pions from breakup of pionium

T.A. Heim[1], **K. Hencken**[1], **M. Schumann**[1], **D. Trautmann**[1] **and G. Baur**[2]

[1] *Institut für Physik, Universität Basel, Klingelbergstrasse 82, CH–4056, Basel, Switzerland*
[2] *Institut für Kernphysik, Forschungszentrum Jülich, 52425 Jülich, Germany*

The experiment DIRAC requires in its analysis a reliable calculation for the number of pionium atoms produced in the target. At small relative momentum $Q < 3\text{MeV}/c$ the observed momentum distribution shows an excess of $\pi^+\pi^-$ pairs (attributable to 'atomic' pairs) as compared to the momentum distribution obtained from a fit at larger relative momentum where only 'free' pairs contribute [1]. However, extracting the number of pionium atoms in this way requires accurate (momentum) distributions of pions from the breakup of pionium as an input.

In the framework of the semiclassical approximation [2], the distribution of the pions from the breakup, resolved with respect to magnitude and angular direction of the momentum of the out-coming pions, is given by

$$\frac{\mathrm{d}^2\sigma_i}{\mathrm{d}\Omega\,\mathrm{d}p} = \int \mathrm{d}^2 b \left| \sum_{l_f,m_f} a_{fi}(b) Y_{l_f,m_f}(\Omega) \right|^2 ,$$

where $a_{fi}(b)$ denotes the amplitude for the transition from the bound initial state i to the final continuum state with (internal) relative momentum p of the pionium. Here, $p = m_{\text{red}}\dot{r} = \frac{1}{2}(p_+ - p_-)$, with $p_\pm$ denoting the momentum of the two pions in their center-of-mass frame. With the method outlined in [2] we can calculate these transitions into the continuum directly for arbitrary initial states. The azimuthal symmetry with respect to the beam axis reduces the angular distribution to a mere θ-dependence where θ denotes the angle between the beam and the out-coming π^+ (or π^-, since the distribution is symmetric under the reflection $\theta \leftrightarrow \pi - \theta$ in first order of the interaction, i.e., considering only breakup of the pionium due to one-photon exchange and taking into account the nearly perfect z-parity conservation [3,4], which becomes exact in the sudden limit).

We have calculated and analyzed the differential cross sections for breakup of pionium on a Nickel target at projectile energy 5 GeV, for initial states n_is and n_ip with $n_i \leq 4$. We found that when integrated over the momentum, the angular distribution shows the two pions to be preferentially emitted perpendicular to the beam, as expected. This strongly non-isotropic distribution is to be contrasted with the isotropic distribution of the 'free' pairs not stemming from the breakup of pionium.

When integrated over the angles, the momentum distribution for breakup from an initial state with principal quantum number n_i has its peak at $p \approx (1/2n_i)$ MeV/c, with a width of approximately $\Delta p \approx (1/2n_i)$ MeV/c. The peak amplitude scales as n_i^2, as expected. Similar calculations for other initial states (d- and f-states reached by multiple scattering) and other target materials will be carried out. These momentum distributions will then be used in a more accurate extraction of the number of pionium atoms produced in the experiment.

[1] L.G. Afanasyev *et al.*, Phys. Lett. B **338** (1994) 478.
[2] Z. Halabuka *et al.*, Nucl. Phys. B **554** (1999) 86.
[3] T.A. Heim, K. Hencken, D. Trautmann and G. Baur, J. Phys. B **33** (2000) 3583.
[4] T.A. Heim, K. Hencken, D. Trautmann and G. Baur, J. Phys. B **34** (2001) 3763.

Higher order corrections to the bound-bound excitation cross sections of pionium

M. Schumann[1], **T. Heim**[1], **K. Hencken**[1], **D. Trautmann**[1] **and G. Baur**[2]

[1] *Institut für Physik, Universität Basel, Klingelbergstr. 82, CH-4056, Basel, Switzerland*
[2] *Institut für Kernphysik, Forschungszentrum Jülich, 52425 Jülich, Germany*

The total and excitation cross sections for the interaction of pionium with the target atoms are an essential theoretical input for the experiment DIRAC [1]. They are well known in the Born approximation and can be calculated to a very high accuracy including various corrections [2–4]. However, Afanasyev et al. [5] showed that the total cross section in Born approximation overestimates the value in Glauber approximation by up to 14% for $Z = 73$ (Ta). This is a major setback for the prospect of calculating the cross sections to 1% as is required for the success of the experiment. To calculate the excitation cross sections we start with the transition amplitude between an initial state i and a final state f given by

$$a^G_{fi}(b) = \int d^3r\, \psi^*_f(\vec{r})\, [1 - \exp\big(i\chi(b,\vec{r})\big)]\, \psi_i(\vec{r}) \ ,$$

where $\psi_{i,f}(\vec{r})$ are non-relativistic hydrogen-like wave functions and $\chi(b,\vec{r}) = -(e/\hbar v) \int [\Phi(\vec{r}\,'/2) - \Phi(-\vec{r}\,'/2)]dz$ with the screened potential $\Phi(r') = (Ze/r') \sum A_k \exp(-\alpha_k r')$. Unlike in the case of the total cross section, where the closure approximation leads to significant simplifications, the excitation cross sections cannot be reduced to a one-dimensional integral in general. Instead a full numerical integration is required. Since the cross sections in Born approximation are well known, we only need to calculate the higher order corrections. Therefore we evaluate the integral

$$\Delta_{fi} = 2\pi \int b\, db\, [|a^B_{fi}(b)|^2 - |a^G_{fi}(b)|^2] \ ,$$

which converges much faster than the full cross section since the higher order corrections are most important for small impact parameters b. Figure 1 shows the results for a few selected transitions. One can see that even for a Nickel target ($Z = 28$) the higher order corrections are between 3% and 5%, while for large Z they reach up to 30%. Clearly, we need to take these results into consideration to reach the desired accuracy of 1% for the cross sections.



Figure 1: Relative difference between Glauber and Born approximation for the excitation cross sections of 1s–2p, 2p–3d, and 2s–3p transitions as a function of the target charge number Z.

[1] B. Adeva et al., CERN/SPSLC 95–1, SPSLC/P 284, Geneva (1995).
[2] Z. Halabuka, et al., Nucl. Phys. B **554** (1999) 86.
[3] T.A. Heim, K. Hencken, D. Trautmann, and G. Baur, J. Phys. B **33** (2000) 3583.
[4] T.A. Heim, K. Hencken, D. Trautmann, and G. Baur, J. Phys. B **34** (2001) 3763.
[5] L. Afanasyev, A. Tarasov, and O. Voskresenskaya, J. Phys. G **25** (1999) B7.

Calculation of electromagnetic breakup of pionium

T. Heim[1], **K. Hencken**[1], **M. Schumann**[1], **D. Trautmann**[1] **and G. Baur**[2]

[1] *Institut für Physik, Universität Basel, Klingelbergstr. 82, CH-4056, Basel, Switzerland*
[2] *Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany*

The calculation of the electromagnetic excitation and breakup of Pionium in the target is an important input for the analysis of the experiment DIRAC at CERN. The cross section for excitation, as well as, breakup was studied within the semiclassical approximation for the dominant Coulomb part of the interaction ("scalar interaction") in [1]. In addition to coherent excitations, where the atom remains in the ground state, incoherent excitations have to be considered also. Summing over all excited states using the closure approximation one obtains a compact expression in terms of the inelastic scattering function S_{inel} [2]. Here only the dominant Coulomb interaction is considered. The S_{inel} were calculated within a Dirac-Hartree-Fock-Slater calculation. Simpler models like "antiscreening" and the "no-correlation limit" were found to be not adequate for the accuracy needed. The contribution of transverse photons was estimated to be at most 0.4% by making use of the long-wavelength approximation.

Calculation of higher order effects are studied within the Glauber approximation and are discussed in [3]. This leaves us now with the calculation of the "magnetic interaction", that is, the current interaction as the estimate show it to contribute almost 1%. In addition we want to study possible effects due to the relativistic treatment of the interaction between pionium and target [4]. The nonrelativistic Schrödinger wave function for the Pionium states could easily be replaced by the solution of the Klein-Gordon equation, but this gives only corrections of order α^2. Starting from the Klein-Gordon equation, we separate terms coming from the external field A_X, treating them as a perturbation. Using the Feshbach-Villars reduction to separate "large" and "small components", we find the interaction between pionium and target to be

$$ieA_{X0} + \frac{ie}{2m}\left[\partial_0 A_{X0} + \vec{\nabla}\vec{A}_X + \vec{A}_X\vec{\nabla}\right] - \frac{e^2}{2m}\left(A_{X0}^2 - \vec{A}_X^2\right) - \frac{e^2}{m}\Phi_C A_{X0}.$$

Besides the usual interaction terms, two new contributions are found: The interaction quadratically in A is given by

$$A_X^\mu A_{X\mu} = A_{X0}^2 - \vec{A}_X^2 = \frac{1}{\gamma^2}A_0^2,$$

which is suppressed by $1/\gamma^2$. The nonrelativistic treatment only leads to $\vec{A}_X^2$ instead, which is not necessarily small. The second term is a Seagull diagram of the form $\Phi_C A_{X0}$ with Φ_C the Coulomb potential between $\pi^+\pi^-$. The same terms are also found starting from a diagrammatic approach, taking into account that Pionium consists of two equally heavy constituents. Explicit calculations of both the magnetic term, as well as, the Seagull diagram are performed. The Seagull diagram is found to contribute only to order 10^{-5}; the magnetic terms are found to be less than 1%, in agreement with the estimates.

[1] Z. Halabuka, et al., Nucl. Phys. B **554** (1999) 86.
[2] T. A. Heim, K. Hencken, D. Trautmann, and G. Baur, J. Phys. B **33** (2000) 3583.
[3] M. Schumann et al., contribution to this miniproceedings.
[4] T. A. Heim, K. Hencken, D. Trautmann, and G. Baur, J. Phys. B **34**, (2001) 3763.

πK atom: observation and lifetime measurement with DIRAC

A. Lanaro

(on behalf of the DIRAC Collaboration)

CERN, CH-1211, Geneva 23, Switzerland

The detection of πK atoms ($A_{\pi K}$) and the measurement of their lifetime provide a model independent way to determine $|a_{1/2}-a_{3/2}|$, the difference between I=1/2,3/2 S-wave πK scattering lengths. The πK scattering probes the $SU(3)_L \times SU(3)_R$ predictions of Chiral Perturbation Theory (ChPT) with u, d, s, and unequal mass kinematics, whereas the investigation of $\pi\pi$ scattering checks the symmetry breaking part of $SU(2)_L \times SU(2)_R$ with u and d quarks only.

The πK scattering amplitude has been evaluated at next-to-leading order in standard ChPT [1]. On the experimental side, S-wave πK phase shifts have been measured from threshold up to 1.3 GeV, using the scattering of moderately energetic charged kaon beams ($2 < E_K < 13$ GeV). The results for the I=1/2 and I=3/2 S-wave πK scattering lengths are spread over a wide range: $0.13 < a_{1/2} < 0.24$, $-0.13 < a_{3/2} < -0.05$, in units of inverse π mass.

To determine the πK threshold parameters, the DIRAC Collaboration proposes the same experimental approach as in the $\pi\pi$ case, which consists in the detection of πK atoms [2].

$A_{\pi K}$ are detected because they have a sizeable probability to break-up in traversing the target material where they are produced. The ratio of experimentally detected $\pi^{\pm}K^{\mp}$ atomic pairs to the full number of produced atoms, yields a measurement of the break-up probability in a given target, and, thus, of the $A_{\pi K}$ lifetime. The relation between the ground state lifetime and the difference $|a_{1/2}-a_{3/2}|$ is given, in first order, by the Deser formula [3]. Using the ChPT prediction at leading order in Isospin breaking for $a_{1/2}$ and $a_{3/2}|$ [1], the lifetime of the $A_{\pi K}$ ground state is predicted to be $4.8 \cdot 10^{-15}$ s.

The Collaboration envisages to use the present detection apparatus, as well as incoming particle beam and energy (24 GeV/c protons from the CERN PS). Only some new implementations to the particle identification system would have to be considered to allow the simultaneous detection of charged pions and kaons. Additional threshold Cherenkov counters of different type have been proposed [2]. Two identical detectors will employ a heavy gas radiator (Freon 114), suitable to discriminate $\pi^{\pm}$ from $K^{\pm}$ in the kinematic range of accepted $A_{\pi K}$. Another Cherenkov detector will employ Aerogel as radiator to discriminate K^{+} from protons.

In the summer of 2001 a test has been performed using the present Cherenkov detectors filled with SF_6 gas. The discrimination of pions from kaons was investigated by detecting the $\phi(1020) \rightarrow K^{+}K^{-}$ resonance decay mode. During 2002, the Collaboration will also perform tests of Aerogel samples with very low index of refraction (n=1.008), manufactured by the Boreskov Institute of Catalysis at Novosibirsk.

In three years of running time, DIRAC expects to detect a few thousands πK atomic pairs, which should lead to a determination of $|a_{1/2}-a_{3/2}|$ with $10\div 15\%$ accuracy, and to a substantial improvement on the precision of the $\pi\pi$ S-wave scattering length parameter $|a_0 - a_2|$.

[1] V. Bernard, N. Kaiser and U. Meissner, Nucl. Phys. B **357** (1991) 129.

[2] B. Adeva et al., the DIRAC Collaboration, CERN/SPSC 2000-032, 17 Aug 2000.

[3] S.M. Bilenky et al., Sov. J. Nucl. Phys. **10** (1969) 469.

Can one extract $\mathrm{m_s}/\hat{\mathrm{m}}$ from πK atom properties?

H. Sazdjian

Groupe de Physique Théorique, Institut de Physique Nucléaire,
Université Paris XI, F-91406 Orsay Cedex, France
E-mail: sazdjian@ipno.in2p3.fr

The lifetime and the $2p-2s$ level energy splitting of the πK atom are studied. The lifetime is essentially dependent on the combination $(a_0^{1/2}-a_0^{3/2})$ of the scattering lengths of the strong interaction process $\pi K \to \pi K$. This combination is independent, at leading order of chiral perturbation theory, of the quark mass ratio $m_s/\hat{m}$. The measurement of the πK atom lifetime would therefore mainly provide information about higher-order (one-loop level) low energy constants. The energy splitting between the $2p-2s$ levels is essentially dependent on the combination $(2a_0^{1/2}+a_0^{3/2})$ of the scattering lengths. This combination depends, at leading order of chiral perturbation theory, on the quark mass ratio $m_s/\hat{m}$. This ratio is however multiplied by a term proportional to the deviation of the $SU(2)\times SU(2)$ quark condensate from the Gell-Mann–Oakes–Renner value. Therefore, if at the $SU(2)\times SU(2)$ level ($\pi\pi$ scattering amplitude) the chiral perturbation theory is close on experimetal grounds to the standard scheme of Gell-Mann, Oakes and Renner and of Gasser and Leutwyler, then the $m_s/\hat{m}$ dependence of the $2p-2s$ energy splitting would hardly be detectable. On the other hand, if sizable deviations from the standard scheme are already present at the $SU(2)\times SU(2)$ level, then these are amplified in the $2p-2s$ energy splitting by the quark mass ratio and experimental detection of the $m_s/\hat{m}$ dependent term becomes possible. In any event, the measurement of the $2p-2s$ energy splitting in the πK atom would provide an experimental test for the standard scheme of chiral perturbation theory. In general, the measurements of the lifetime and of the energy splitting would allow the determination of each of the scattering lengths $a_0^{1/2}$ and $a_0^{3/2}$.

πK scattering and dispersion relations

B. Ananthanarayan[1], **P. Büttiker**[2] **and B. Moussallam**[3]

[1] *Centre for Theoretical Studies, Indian Institute of Science, Bangalore 560 012, India*
[2] *Institut für Kernphysik, Forschungszentrum Jülich, 52425 Jüelich, Germany*
[3] *Groupe de Physique Théorique, IPN, Université Paris–Sud, 91406 Orsay Cédex, France*

Chiral perturbation theory provides the low energy effective theory of the standard model that describes the interactions involving hadronic degrees of freedom. As a non–renormalizable theory, it requires additional low energy constants (LEC) that have to be introduced at each order of the loop– or momentum–expansion. In $SU(3)$ at next–to–leading order, ten LECs $L_i(\mu)$ enter the calculations. However, these quantities cannot be fixed by chiral symmetry only; experimental information is needed to pin them down. Using the K_{l4} form factors and $\pi\pi$ sum rules provide accurate estimates for some of the LECs (e.g. L_1, L_2, L_3) but leave the large N_c–suppressed LEC L_4 essentially undetermined. This is due to the fact that in such processes, contributions from L_4 accidentally are suppressed. Such a suppression does not occur in πK scattering [1]. Therefore this process is suitable for a more reliable determination of L_4 and also provides estimates for L_1, L_2 and L_3. We use dispersion relations to fix these quantities [2], relying on analyticity, unitarity and crossing symmetry only, which are the suitable tools for a comparison of experimental data with ChPT. By combining fixed–t and hyperbolic dispersion relations for $T^+(s,t,u)$ and $T^-(s,t,u)$ and rewriting the chiral amplitudes, we find an appropriate matching of the chiral and the dispersive representations of these amplitudes. Saturating the dispersive integrals with the available data [3], we find the following estimates

$$\begin{aligned} L_1^r(m_\rho) &= 0.84 \pm 0.15, & L_2^r(m_\rho) &= 1.36 \pm 0.13, \\ L_3^r(m_\rho) &= -3.65 \pm 0.45, & L_4^r(m_\rho) &= 0.22 \pm 0.3, \end{aligned}$$

yielding an estimate for L_4 with improved reliability.

[1] V. Bernard, N. Kaiser and U.–G. Meißner, Phys. Rev. D **43**, 2757 (1991); Nucl. Phys. B **357**, 129 (1991).

[2] B. Ananthanarayan and P. Büttiker, Eur. Phys. J. C **19**, 517 (2001) [hep-ph/0012023]; B. Ananthanarayan, P. Büttiker and B. Moussallam, Eur. Phys. J. C (2001), to appear.

[3] P. Estabrooks et al., Nucl. Phys. B **133**, 490 (1978); D. Aston et al., Nucl. Phys. B **296**, 493 (1988); D. Cohen et al., Phys. Rev. D **22**, 2595 (1988); A. Etkin et al., Phys. Rev. D **25**, 1786 (1982)

Isospin violation in pion–kaon scattering

B. Kubis and U.-G. Meißner

Forschungszentrum Jülich, Institut für Kernphysik (Theorie), D–52425 Jülich, Germany

Pion–kaon scattering near threshold is one of the cleanest processes to test our understanding of chiral dynamics in the presence of strange quarks. The existing determinations of the S–wave scattering lengths are, however, plagued by large uncertainties, such that high hope lies in the extraction of these parameters from πK bound state experiments [1].

In order to relate the lifetime of $\pi^- K^+$ (or $\pi^+ K^-$) atoms to one particular combination of the pion–kaon scattering lengths, one has to make use of a modified Deser formula for the decay into the neutral $\pi^0 K^0$ channel which includes next–to–leading order effects in isospin breaking,

$$\Gamma_{\pi^0 K^0} \propto \left(a_0^{3/2} - a_0^{1/2} + \epsilon\right)^2 (1 + K) \tag{1}$$

(see [2] for an equivalent formula for the case of pionium). Here, ϵ represents the isospin violating correction in the scattering amplitude $\pi^- K^+ \to \pi^0 K^0$ at threshold, while K is an additional contribution only calculable within the bound state formalism.

We have evaluated the parameter ϵ up to one–loop order, including leading effects both in the quark mass difference $m_u - m_d$ and the electromagnetic fine structure constant $\alpha = e^2/4\pi$ [3] (see also [4]). While $m_u - m_d$ effects are forbidden at leading order for $\pi\pi$ scattering and pionium and thus numerically negligible in comparison to electromagnetic effects, there are contributions linear in $m_u - m_d$ to the πK scattering lengths which are of comparable size to the corrections of electromagnetic origin. At one–loop order, virtual photon loops induce a kinematical divergence at threshold (the Coulomb pole) $\propto 1/|\mathbf{q}_{\rm in}|$ which has to be subtracted in order to properly define the scattering length. Furthermore, photon loops contain an infrared divergence, which however vanishes at threshold. The corresponding radiative process $\pi^- K^+ \to \pi^0 K^0 \gamma$ which has to be included in order to define an infrared finite total cross section yields a totally negligible contribution at threshold.

The combined result for the scattering length, expressed in terms of relative corrections to the tree level isospin symmetric one, numerically reads

$$\begin{aligned} a_0(\pi^- K^+ \to \pi^0 K^0) &= \frac{\sqrt{2}}{3}\left(a_0^{3/2} - a_0^{1/2}\right)^{\rm tree} \Big\{ 1 + \underbrace{0.006}_{\mathcal{O}(m_u - m_d)} + \underbrace{0.009}_{\mathcal{O}(e^2)} \\ &\quad + \underbrace{(0.121 \pm 0.009)}_{\mathcal{O}(p^4)} + \underbrace{(0.008 \pm 0.002)}_{\mathcal{O}(p^2(m_u - m_d))} - \underbrace{(0.009 \pm 0.008)}_{\mathcal{O}(p^2 e^2)} \Big\} . \end{aligned} \tag{2}$$

The error ranges displayed for the next–to–leading order contributions are due to the uncertainties in the low–energy constants entering at that order. The value for the lifetime of the πK atom obtained using the "uncorrected" Deser formula, $\tau_{\pi K} = (4.8 \pm 0.1) \times 10^{-15}$ s, is slightly reduced to $\tau_{\pi K} = (4.7 \pm 0.2) \times 10^{-15}$ s. The additional correction K in (1) however remains to be calculated.

[1] B. Adeva et al., CERN-SPSC-2000-032, CERN-SPSC-P-284-ADD-1, Aug 2000.

[2] A. Gall et al., Phys. Lett. B **462** (1999) 335; J. Gasser et al., Phys. Lett. B **471** (1999) 244.

[3] B. Kubis and U.-G. Meißner, arXiv:hep-ph/0107199, Nucl. Phys. A, in print.

[4] A. Nehme and P. Talavera, arXiv:hep-ph/0107299.

The first measurement of kaonic nitrogen with DEAR at DAΦNE

C. Curceanu (Petrascu)[1]
on behalf of the DEAR Collaboration

[1] *LNF-INFN, Via E. Fermi 40, 00044 Frascati (Roma), Italy*
(on leave from IFIN-HH, Bucharest, Romania)

In May 2001 the DEAR (DAΦNE Exotic Atom Research) collaboration [1] performed the first measurement of an exotic atom (kaonic nitrogen) at the DAΦNE collider of the Laboratori Nazionali di Frascati dell'INFN. The measurement was performed with a cryogenic target (118 K) filled with nitrogen at about 2 bar ($\rho \simeq 4.5\rho_{NTP}$). As detector, 8 CCD-22 were used. A total integrated luminosity of about 2 pb^{-1} was collected. After an optimization of the machine performances and of the DEAR degrader, an X-ray spectrum was obtained and analysed by performing a global fit. The global fit (with a χ^2/NDOF = 0.95) gave a statistical significance of the two kaonic nitrogen lines measurable with DEAR of about 3σ for the line at 4.58 keV ($7 \rightarrow 6$ transition) and of about 3.7σ for the line at 7.6 keV ($6 \rightarrow 5$ transition), giving a global statistical significance of about 5σ. These results compare favourable with the Monte Carlo simulation results with an yield of the two transitions of about $90^{+10}_{-25}\%$. An empty target-like measurement was as well performed. The background subtracted spectrum is presented in Figure 1. The experiment could therefore demonstrate the capability of creating and detecting exotic atoms using the kaons flux from DAΦNE.

Next stage of the experiment is the measurement of the K-complex in kaonic hydrogen.



Figure 1: The kaonic nitrogen background subtracted spectrum. X-ray lines are: 1 = Al K$_\alpha$; 2 = Si K$_\alpha$; 3 = Sc K$_\alpha$; 4 = **KN (4.58 keV)**; 5 = V K$_\alpha$; 6 = Cr K$_\alpha$; 7 = Cr K$_\beta$ + Mn K$_\alpha$; 8 = Fe K$_\alpha$ + Mn K$_\beta$; 9 = Fe K$_\beta$; 10 = **KN (7.62 keV)**; 11 = Cu K$_\alpha$; 12 = Zn K$_\alpha$; 13 = Zn K$_\beta$; 14 = Pb L$_\alpha$

[1] S. Bianco *et al.*, Rivista del Nuovo Cimento **22**, No. 11 (1999) 1.

The quark condensate on the lattice

P. Hasenfratz

Institut für Theoretische Physik, Universität Bern, Sidlerstr. 5, 3012 Bern, Switzerland

Chiral symmetry in QCD with $N_f \geq 2$ massless quark flavours is expected to be broken spontaneously producing a non-zero $\langle\bar{\psi}\psi\rangle$ quark condensate generated through non-perturbative effects. A direct study of this condensate requires chiral symmetric lattice fermions (domain-wall, overlap, or the fixed-point formulations) which are expensive to simulate. The fermion condensate

$$\langle\bar{\psi}\psi\rangle = -\lim_{m_q\to 0}\lim_{V\to\infty}\frac{d}{dm_q}\log Z \qquad (1)$$

is related to the low energy constant Σ of the tree level effective chiral lagrangean as $\langle\bar{\psi}\psi\rangle = -N_f\Sigma$. A way to determine the low energy constant Σ is to measure the condensate in a small volume V in a fixed topological sector Q of QCD: $N_f\Sigma_{m_q,V,Q} = -\langle\bar{\psi}\psi\rangle_{m_q,V,Q}$ and use chiral perturbation theory to connect it to Σ defined at $m_q = 0$ and $V = \infty$. Until now this is carried through in the quenched approximation only. Chiral perturbation theory predicts in this case [1]

$$\lim_{m_q\to 0}\frac{\Sigma_{m_q,V,Q} - |Q|/m_qV}{m_q} = \frac{\Sigma^2 V}{2|Q|} + \frac{c}{a^2}\,, \qquad (2)$$

where $|Q|/m_qV$ is the contribution from the exact zero modes, while c/a^2 (a is the lattice unit) is the ultraviolet divergent cut-off effect induced by the finite quark mass. Measuring $\Sigma_{m_q,V,Q}$ in different volumes, eq. (2) allows to find Σ. The bare low energy constant Σ is multiplicatively renormalized. The renormalization factor between Σ and $\Sigma_{\overline{MS}}(\mu)$ is calculated by combining non-perturbative measurements and continuum perturbation theory. We obtained the result $\Sigma_{\overline{MS}}(2\mathrm{GeV}) = (270 \pm 10\mathrm{MeV})^3$ [2], which, using the GMOR relation, corresponds to $\overline{m}_{\overline{MS}}(2\mathrm{GeV}) = (3.9 \pm 0.4)\mathrm{MeV}$ for the averaged u,d quark mass. This result is obtained at a resolution $a \approx 0.13$fm and no systematic continuum extrapolation is performed yet. The same applies also to the other measurements (they are consistent with our result) available at this moment [3]. One should keep in mind that these results refer to the quenched approximation ($N_f = 0$). It is unknown how far is the $N_f = 0$ low energy constant from that at $N_f = 3$. The number above (which is close to the phenomenologically expected one) suggests that $\Sigma(N_f)$ depends smoothly on N_f. This is not true, however for the condensate $\langle\bar{\psi}\psi\rangle$ itself. The $N_f = 0$ (quenched) QCD is not a healthy QFT and the tree level relation $\langle\bar{\psi}\psi\rangle = -N_f\Sigma$, unlike in full QCD, receives corrections which are even diverging in the chiral limit. As opposed to the low energy constant Σ, the condensate is not well defined in the quenched approximation [1].

[1] J.C. Osborn et al., Nucl. Phys. B540 (1999) 317; P.M. Damgaard et al., Nucl. Phys. B547 (1999) 305.

[2] P. Hasenfratz, S. Hauswirth, K. Holland, T. Jörg and F. Niedermayer, arXiv:hep-lat/0109007.

[3] P. Hernandez et al., JHEP 0107 (2001) 018; T. DeGrand, arXiv:hep-lat/0107014; L. Giusti et al., arXiv:hep-lat/0108007.

Chiral and lattice calculations of the $\pi\pi$ scattering lengths

Gilberto Colangelo

Institut für Theoretische Physik der Universität Zürich Winterthurerstr. 190 CH-8057 Zürich

In this talk I have reviewed the status of lattice calculations of the pion scattering lengths from a nonexpert point of view. The history of such calculations starts in the eighties/early nineties. I have split the discussion by reviewing first the calculations pre-2000 and then those post-2000. The most advanced calculation of last century is that of Fukugita et al. [1], where reference to earlier literature can also be found. This paper is also particularly relevant because, as far as I know, it is the only attempt at calculating the scattering length in the $I = 0$ channel on the lattice. Despite the huge effort and the remarkable results, various systematic effects were not under good control.

A new generation of calculations by two japanese collaborations have started in recent years [2,3]. Although the final results are not yet available, encouraging and very interesting preliminary results for the $I = 2$ scattering lengths have been presented at recent Lattice conferences.

I have also discussed the recent results of a small chinese collaboration that has exploited the use of improved actions to do the calculation on a very coarse and anisotropic lattice [4], and therefore perform it on a relatively small computer. I sincerely hope that this brave and encouraging calculation will challenge also smaller lattice groups to attempt such nontrivial but extremely interesting calculation.

The agreement between experiments, CHPT, and lattice calculations for the $\pi\pi$ scattering lengths would represent a tremendous achievement for our understanding of strong interactions The CHPT calculation has reached a very high level of accuracy [5], and recent experimental results have confirmed the prediction [6]. Hopefully, the lattice calculations will also reach the same level of accuracy one day.

[1] M. Fukugita, Y. Kuramashi, M. Okawa, H. Mino and A. Ukawa, Phys. Rev. D **52** (1995) 3003 [arXiv:hep-lat/9501024].

[2] S. Aoki *et al.* [JLQCD Collaboration], Nucl. Phys. Proc. Suppl. **83** (2000) 241 [arXiv:hep-lat/9911025].

[3] S. Aoki *et al.* [CP-PACS Collaboration], arXiv:hep-lat/0110151.

[4] C. Liu, J.h. Zhang, Y. Chen and J.P. Ma, arXiv:hep-lat/0109020; arXiv:hep-lat/0109010.

[5] G. Colangelo, J. Gasser and H. Leutwyler, Phys. Lett. B **488** (2000) 261 [arXiv:hep-ph/0007112].

[6] S. Pislak *et al.* [BNL-E865 Collaboration], Phys. Rev. Lett. **87** (2001) 221801 [arXiv:hep-ex/0106071].

The quark condensate from K_{e4} decays

G. Colangelo[1], **J. Gasser**[2] **and H. Leutwyler**[2]

[1] Institute for Theoretical Physics, University of Zürich
Winterthurerstr. 190, CH-8057 Zürich, Switzerland
[2] Institute for Theoretical Physics, University of Bern
Sidlerstr. 5, CH-3012 Bern, Switzerland

Roy equations [1] combined with chiral symmetry allow one to predict [2] the low-energy behaviour of the $\pi\pi$ scattering amplitude with high precision. The prediction is based on the hypothesis that the quark condensate is the leading order parameter of spontaneously broken chiral symmetry. This has been questioned in Refs. [3,4], where it is emphasized that experimental evidence for this scenario is not available. The recent result from the high statistics K_{e4} experiment E865 at Brookhaven [5] closes this gap, because those data allow one to determine the size of the leading term in the quark mass expansion of the pion mass. E865 confirms [5] the hypothesis that underlies our prediction of the $\pi\pi$ S-wave scattering lengths: more than 94% of the pion mass stems from the quark condensate. The generalized framework of $SU(2)\times SU(2)$ chiral perturbation theory developed in [3], that allows for a small or vanishing condensate, can therefore be dismissed.

Note added: In Ref. [6], the E865 data [5] have been analyzed without the use of chiral symmetry constraints.

[1] S.M. Roy, Phys. Lett. B **36** (1971) 353.

[2] G. Colangelo, J. Gasser and H. Leutwyler, Phys. Lett. B **488** (2000) 261 [hep-ph/0007112]; Nucl. Phys. B **603** (2001) 125 [hep-ph/0103088]; Phys. Rev. Lett. **86**, 5008 (2001) [hep-ph/0103063].

[3] M. Knecht, B. Moussallam, J. Stern and N.H. Fuchs, Nucl. Phys. B **457** (1995) 513 [hep-ph/9507319]; *ibid.* B **471** (1996) 445 [hep-ph/9512404].

[4] L. Girlanda, M. Knecht, B. Moussallam and J. Stern, Phys. Lett. B **409** (1997) 461 [hep-ph/9703448].

[5] S. Pislak *et al.* [BNL-E865 Collaboration], Phys. Rev. Lett. **87** (2001) 221801 [hep-ex/0106071].

[6] S. Descotes, N.H. Fuchs, L. Girlanda and J. Stern, arXiv:hep-ph/0112088.

Radiative corrections to semileptonic decays

M. Knecht

Centre de Physique Théorique, CNRS Luminy Case 907, F-13288 Marseille Cedex 9

Semileptonic decays of the light pseudoscalar mesons provide important informations on the non-perturbative aspects of the QCD vacuum (low-energy $\pi\pi$ scattering and low-energy constants L_i from $K_{\ell 4}$ decay), or on the CKM matrix elements V_{ud} (pion β decay) and V_{us} ($K_{\ell 3}$ decay). An accurate determination of these quantities requires high statistics experiments, as well as reliable theoretical descriptions. In particular, isospin breaking effects, arising both from $m_u \neq m_d$ and from radiative corrections, should be included. So far, this has not been done in a systematic way.

A general framework allowing for a systematic treatment of isospin breaking effects within the chiral low-energy expansion has been introduced in [1], and applied to the simplest, $\pi_{\ell 2}$ and $K_{\ell 2}$ semileptonic modes. Recently, the $K_{\ell 3}$ channels have been treated as well [2] (for a preliminary discussion of the $K_{\ell 4}$ modes, see [3]).

In the presence of radiative corrections, the factorization between the hadronic and leptonic currents in semileptonic decays is lost. In the case of the $K_{\ell 3}$ transitions, this entails that the two form factors $f_{\pm}$ not only depend on the momentum transfer between the two mesons, but also on a second, independent, kinematical variable. Furthermore, the necessity to include the decay mode with a soft photon emission, in order to obtain an infrared finite result, modifies the structure of the Dalitz plot density. Explicit expressions for the $K_{\ell 3}$ form factors at one loop with isospin breaking effects included have been obtained in [2], where existing expressions [4] for the Dalitz plot distribution have been corrected for several mistakes.

The effects of isospin breaking on the extraction of V_{us} along the lines described in [5] have also been discussed and found to be small (see [2] for details). A reliable estimate of the two-loop strong interaction effects appears more than ever as being the crucial issue for a precise determination of V_{us} from $K_{\ell 3}$ decay [6].

[1] M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, Eur. Phys. J. C **12** (2000) 469 [hep-ph/9909284].

[2] V. Cirigliano, M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, submitted to Eur. Phys. J. C [hep-ph/0110153].

[3] V. Cuplov, contribution to this meeting.

[4] E.S. Ginsberg, Phys. Rev. **142** (1966) 1035; **162** (1967) 1570 [Erratum-ibid. **187** (1967) 2280]; **171** (1968) 1675 [Erratum-ibid. **174** (1968) 2169]; Phys. Rev. D **1** (1970) 229.

[5] H. Leutwyler and M. Roos, Z. Phys. C **25** (1984) 91.

[6] N.H. Fuchs, M. Knecht and J. Stern, Phys. Rev. D **62** (2000) 033003 [hep-ph/0001188].

Isospin violation in $K_{\ell 4}$ decays

Vesna Cuplov

Centre de Physique Théorique, CNRS Luminy, Case 907, F-13288 Marseille Cedex 9, France

The framework used in Ref.[1] for studying $K_{\ell 4}$ decays is modified in order to account for isospin violating contributions, $m_u \neq m_d$ and $\alpha \neq 0$. The full kinematics, which requires three reference frames and five variables introduced by Cabibbo and Maksymowicz Ref.[2] is changed. It is necessary to introduce a tensorial form factor, $T_{\mu\nu} = \frac{1}{M_K^2} T p_{1\mu} p_{2\nu}$, in the matrix element:

$$\mathcal{T} \;=\; \frac{G_F V_{us}^{\star}}{\sqrt{2}} \left[\; \bar{u}(p_\nu)\gamma_\mu(1-\gamma_5)v(p_l)(V^\mu - A^\mu) + \;\bar{u}(p_\nu)\sigma_{\mu\nu}(1+\gamma_5)v(p_l)T^{\mu\nu} \;\right]$$

However this tensorial form factor appears only in the $\pi^+\pi^-$ channel, due to electromagnetic corrections.

The decay rate is $d\Gamma_5 = G_F^2 \mid V_{us} \mid^2 N(s_\pi, s_l) J_5(s_\pi, s_l, \theta_\pi, \theta_l, \phi) \times ds_\pi ds_l d(\cos\theta_\pi) d(\cos\theta_l) d\phi$. We have a general formula for J_5 written in terms of the mass of the kaon which decays, the masses of the two pions in the final state and all the possible form factors, including the tensorial one. We explicitly calculate the form factors for $K_{\ell 4}$ at tree level using an effective Lagrangian, where not only the pseudoscalars but also the photon and the light leptons have been included as dynamical degrees of freedom Ref.[3].

The numerical values obtained for the decay rates(MeV) and phase space(MeV^3) are shown in the table below. We compare to the values obtained by Bijnens and al. Ref.[4].

Decay	(1) $\pi^+\pi^-e^+\nu_e$	$\pi^+\pi^-\mu^+\nu_\mu$	(2) $\pi^o\pi^o e^+\nu_e$	$\pi^o\pi^o\mu^+\nu_\mu$	(3) $\pi^o\pi^- e^+\nu_e$	$\pi^o\pi^-\mu^+\nu_\mu$
Ref.[4] Tree level	1297	155	683	102	561	55
Masses as in Ref.[4] $m_u = m_d$ $e = 0$ $T = 0$	1292	154	681	101	561	55
phase space MeV^3	1.140	0.146	0.674	0.099	1.339	0.197
Physical masses $m_u \neq m_d$ $e \neq 0$ $T \neq 0$	1292	153	730	110	543	51
phase space MeV^3	1.140	0.146	0.674	0.099	1.307	0.187

In Ref.[4], the masses of the particles are (M_π, M_K)=(139.6, 493.6) MeV, (135, 493.6) MeV and (137, 497.7) MeV for the decays (1), (2) and (3), respectively.

In case of no isospin violation, our Fortran routine reproduces the results of Bijnens and al. at better than 0.5 %. The interpretation of these numerical values leads to the result that the isospin violation, at tree level, is due to form factors for the channel $\pi^o\pi^o$ and to phase space for the channel $\pi^o\pi^-$. The channel $\pi^+\pi^-$ is the only one which involves the tensorial form factor. At tree level, the effects of isospin violation and the existence of a tensorial form factor are numerically tiny in this channel. The calculations at one loop level are in progress.

[1] The second DaΦne physics handbook - Vol.1.

[2] N. Cabibbo and A. Maksymovicz, Phys. Rev. **137** (1965) B438; erratum Phys. Rev. **168** (1968) 1926.

[3] M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, Eur. Phys. J. C **12** (2000) 469.

[4] J. Bijnens, G. Colangelo and J. Gasser, Nucl. Phys. B **427** (1994) 427.

Resonance estimates of low-energy constants in chiral Lagrangians and QCD short-distance constraints

M. Knecht and A. Nyffeler

Centre de Physique Théorique, CNRS-Luminy, Case 907, F-13288 Marseille Cedex 9, France

At higher orders in the chiral expansion and when radiative corrections are included, a large number of unknown low-energy constants enters in ChPT. Lacking enough data, one often relies on estimates for these constants using some resonance Lagrangian. This approach was quite successful for the L_i at order p^4 [1], although one has to make sure that QCD short-distance constraints are met [2]. Starting from the study of the low-energy and high-energy behaviors of the QCD three-point functions $\langle VAP\rangle$, $\langle VVP\rangle$ and $\langle AAP\rangle$, we have evaluated in [3] several $\mathcal{O}(p^6)$ low-energy constants of the chiral Lagrangian within the framework of the lowest meson dominance (LMD) approximation to the large-N_C limit of QCD. In certain cases, values that differ substantially from estimates based on a resonance Lagrangian are obtained:

	C_{78}	C_{82}	C_{87}	C_{88}	C_{89}	C_{90}
LMD	1.09	-0.36	0.40	-0.52	1.97	0.0
Set I	1.09	-0.29	0.47	-0.16	2.29	0.33
Set II	1.49	-0.39	0.65	-0.14	3.22	0.51

	A_{V,p^2}	$A_{V,(p+q)^2}$	A_{A,p^2}	$A_{A,(p+q)^2}$
LMD	-1.11	-0.26	-0.14	0.040
Set I	-1.13	0.0	-0.096	0.0

In the table we have collected our estimates (LMD) for the low-energy constants C_i (even parity) from [4] and A_i (odd parity) from [5], all in units of $10^{-4}/F_0^2$, and compared them to those obtained with a resonance Lagrangian, using different sets of parameters in the Lagrangian (set I and II). We have introduced the abbreviations $A_{V,p^2} = A_2 - 4A_3$, $A_{V,(p+q)^2} = A_2 - 2A_3 - 4A_4$, $A_{A,p^2} = A_{11} + A_{23} - 4A_{24}$, and $A_{A,(p+q)^2} = 3A_{11} + 4A_{17} + A_{23} - 2A_{24} - 4A_{25}$.

The differences, in particular in $C_{88}, C_{90}, A_{V,(p+q)^2}$, and $A_{A,(p+q)^2}$, arise through the fact that QCD short-distance constraints are in general not correctly taken into account in the approaches using resonance Lagrangians. Whereas in the case of the vector form factor of the pion our result for a certain combination of low-energy constants at order p^6 agrees with the one [6] from the resonance Lagrangian, our estimate for one of the counterterm contributions in the form factor A in the decay $\pi \to e\nu_e\gamma$ is by a factor of 5 smaller than the resonance estimate in [7].

Finally, in Ref. [3] we also briefly discussed how to systematically improve our approach. For instance, for the pion-photon-photon transition form factor it is necessary to include a second vector resonance (LMD+V) in order to correctly reproduce the CLEO data [8] for the form factor at high Q^2, with one photon on-shell. In this way one obtains the new estimate $A_{V,p^2}^{LMD+V} = -1.36 \times 10^{-4}/F_0^2$, i.e., about 20% away from the LMD estimate given in the table.

[1] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B **321** (1989) 311.

[2] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, Phys. Lett. B **223** (1989) 425.

[3] M. Knecht and A. Nyffeler, Eur. Phys. J. C **21** (2001) 659.

[4] J. Bijnens, G. Colangelo and G. Ecker, JHEP **9902** (1999) 020; [hep-ph/9902437].

[5] H.W. Fearing and S. Scherer, Phys. Rev. D **53** (1996) 315.

[6] J. Bijnens, G. Colangelo and P. Talavera, JHEP **9805** (1998) 014; [hep-ph/9805389].

[7] J. Bijnens and P. Talavera, Nucl. Phys. B **489** (1997) 387.

[8] J. Gronberg *et al.* [CLEO Collaboration], Phys. Rev. D **57** (1998) 33.

The new pionic hydrogen experiment at PSI

M. Hennebach for the πH collaboration (www.pihydrogen.psi.ch)

Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich

A new high–precision determination of the strong–interaction shift (ϵ_{1s}) and broadening (Γ_{1s}) of the ground state in pionic hydrogen (πH) has been started at PSI [1]. This constitutes a direct measurement of the πN scattering lengths a^+ and a^- and is an important test of the methods of chiral perturbation theory.

The experimental approach is to determine the energies of x-rays emitted by radiative transitions to the 1s–state in pionic hydrogen. This state is both shifted in energy because of the strong interaction between proton and pion and broadened because its lifetime is shortened due to the additional decay channels. The energy of the *np–1s* transitions in πH is in the range of a few keV and should be determined with an accuracy in the 20 meV range. This can be achieved only with a crystal spectrometer.

A previous experiment at PSI [2] measured the *3p–1s* transition in pionic hydrogen and deuterium (πD), combining a highly efficient stopping arrangement (cyclotron trap) for the pion beam and a cylindrically bent Bragg crystal with newly developed CCD (Charge Coupled Device) detectors. While an accuracy of 0.5% was reached for ϵ_{1s}, the error for Γ_{1s} (8%) was an order of magnitude higher, which makes it hard to determine both a^+ and a^- from the πH measurement alone. The πD shift measurement can be used to improve the situation, but this necessitates a good theoretical understanding of the 3–body–problem in the deuterium system.

The goal of this experiment is to reduce the uncertainty of Γ_{1s} to $\approx$1% and that of ϵ_{1s} to 0.2%. This can be achieved by using an improved cyclotron trap, spherically bent crystals and a new large–area CCD for a higher x-ray rate and better background suppression.

During the first production run the *3p-1s* transition was measured at three different pressures (4 bar, 28 bar and liquid hydrogen) to investigate the possible influence of mechanisms that depend on the collision rate between πH atoms and target molecules. One process is the formation of molecular states ($\pi^-\mathrm{H} + \mathrm{H}_2 \rightarrow [(\pi^-\mathrm{pp})\mathrm{p}]_{\mathrm{ee}}$) that could skew the ϵ_{1s} measurement. The other is Coulomb deexcitation ($\pi^-\mathrm{H}_{(\mathrm{n})} + \mathrm{H}_2 \rightarrow \pi^-\mathrm{H}_{(\mathrm{n}')} + \mathrm{H}_2$), a non–radiative process that leads to a Doppler broadening of the line.

A preliminary analysis of the data (Table 1) does not show a pressure dependence of the shift ϵ_{1s}. Furthermore, the averaged result does not deviate from the earlier result [2] which gave $\epsilon_{1s} = 7108 \pm 13 \pm 34$ at 15 bar. The width Γ_{1s} (including the Coulomb deexcitation) appears to be pressure dependent. This will be investigated in the next stage of the experiment, starting in Spring 2002.

Target pressure	ϵ_{1s} [meV]	Γ_{1s} [meV]
4 bar	$7082 \pm 31 \pm 15$	$973 \pm 75 \pm 10$
28 bar	$7137 \pm 18 \pm 40$	$969 \pm 26 \pm 10$
liquid (∼700 bar)	$7095 \pm 25 \pm 25$	$1052 \pm 58 \pm 10$

Table 1: Preliminary fit results for the Summer 2001 data (value ± stat. ± sys.)

[1] Proposal R-98-01.1 at PSI.

[2] H.-Ch. Schröder et al., Eur. Phys. J. **C 21** (2001) 433 vol. 3

Atomic cascade in hadronic atoms

T.S. Jensen[1,2] **and V.E. Markushin**[1]

[1] *Paul Scherrer Institute, CH-5232, Villigen PSI, Switzerland*
[2] *Institut für Theoretische Physik der Universität Zürich,*
Winterthurerstrasse 190,
CH-8057 Zürich, Switzerland

The atomic cascades in exotic hydrogen atoms x^-p ($x^- = \mu^-, \pi^-, K^-, \bar{p}$) have been studied using a new Monte Carlo code that calculates the evolution of the x^-p kinetic energy distribution from the very beginning of the cascade. The cascade model is based on new cross sections for the collisional processes: Stark mixing, elastic scattering, nuclear absorption during collisions, Coulomb deexcitation, and external Auger effect. For the final part of the cascade ($n \leq 5$), Stark mixing, deceleration, and nuclear absorption during collisions for x^-p scattering from hydrogen atoms have been computed in the close–coupling approximation [1-3]; this fully quantum mechanical framework takes threshold and strong absorption effects into account. For $n \geq 8$, a classical–trajectory Monte Carlo model [3] for exotic atoms scattering from molecular hydrogen has been used to calculate the cross sections for Stark mixing, elastic scattering, and Coulomb deexcitation. The semiclassical (eikonal) approximation is used for values of n between the quantum mechanical and classical domains.

The predictions of the detailed kinetics cascade model have been confronted with the experimental observables: X-ray yields, cascade times, and kinetic energy distributions [4]. The case of pionic hydrogen is of particular interest. The new PSI experiment [5], whose goal is to determine the strong interaction width Γ_{1S} of pionic hydrogen on the level of 1%, relies crucially on a good understanding of the kinetic energy evolution during the atomic cascade because the measured $nP \rightarrow 1S$ line profiles must be corrected for the Doppler broadening (see Fig. 1).



Figure 1: The calculated energy profile of the K_β line at 3 bar in pionic hydrogen with Doppler broadening taken into account (solid line) in comparison with the natural line shape assuming $\Gamma_{1S} = 0.87$ eV (dashed line).

[1] T.S. Jensen and V.E. Markushin, PSI-PR-99-32 (1999), arXiv:nucl-th/0001009.

[2] T.S. Jensen and V.E. Markushin, Nucl. Phys. A **689** (2001) 537.

[3] T.S. Jensen and V.E. Markushin, Hyperfine Interactions, in press.

[4] V.E. Markushin and T.S. Jensen, Hyperfine Interactions, in press.

[5] D. Gotta, πN Newsletter **15** (1999) 276. See also M. Hennebach, these proceedings.

Pion-nucleon analysis

M.E. Sainio

Helsinki Institute of Physics, and Department of Physical Sciences, University of Helsinki,
P.O. Box 64, FIN-00014 Helsinki, Finland

There is information of the pion-nucleon interaction in different charge channels available at low energy from a variety of experiments: 1) cross section information, 2) analyzing powers, 3) forward D^+ amplitude, 4) pionic hydrogen level shift and width measurements. Over the years such information has been analysed with different techniques, e.g.

1. Karlsruhe fixed-t analysis KH78/80 [1], partial wave dispersion relation analysis KA.84 [2] and partial wave relation analysis KA.85 [3].
2. VPI-GWU analysis with fixed-t constraints, accessible through the SAID facility `http://gwdac.phys.gwu.edu/`
3. Gashi et al. analysis at very low energies with an effective range expansion [4].
4. Low-energy analysis exploiting forward dispersion relations [5].

A major problem in extracting hadronic quantities from observables is the question of the treatment of the electromagnetic effects. In low-energy scattering the formalism of Tromborg et al. [6] is often used, but there are effects not taken into account by that method. Also, for pionic hydrogen, determining the hadronic scattering lengths requires a careful analysis of electromagnetic corrections [7,8]. For the low-energy scattering region a $\mathcal{O}(p^3)$ calculation in chiral perturbation theory including all electromagnetic effects to that order has recently been published [9].

It turns out, especially in the analysis of sensitive quantities like the Σ-term, that in addition to the electromagnetic corrections also the high partial waves are important. Therefore, it is well motivated to start a new analysis in the spirit of Karlsruhe [10].

[1] R. Koch and E. Pietarinen, Nucl. Phys. A **336** (1980) 331.
[2] R. Koch, Z. Phys. C **29** (1985) 597.
[3] R. Koch, Nucl. Phys. A **448** (1986) 707.
[4] A. Gashi et al., arXiv:hep-ph/0009081.
[5] J. Gasser et al., Phys. Lett. B **213** (1988) 85.
[6] B. Tromborg at al., Phys. Rev. D **15** (1977) 725.
[7] D. Sigg et al., Nucl. Phys. A **609** (1996) 310.
[8] V.E. Lyubovitskij and A. Rusetsky, Phys. Lett. B **494** (2000) 9.
[9] N. Fettes and U.-G. Meißner, Nucl. Phys. A **693** (2001) 693.
[10] P. Piirola et al., arXiv:hep-ph/0110044.

Theory of low-energy pion-nucleon scattering

Nadia Fettes and Ulf-G. Meißner

Forschungszentrum Jülich, Institut für Kernphysik (Theorie), D-52425 Jülich, Germany

In the Standard Model there are two sources of isospin violation. Strong isospin violation is driven by the quark mass difference $(m_u - m_d)$ whereas electromagnetic (em) isospin violation is due to the different quark charges. To extract the strong contribution, which leads to small effects of the size $(m_u - m_d)/\Lambda_{\rm QCD} \sim 2\%$, one thus has to be able to simultaneously treat both sources. In the two flavor sector of QCD, pion-nucleon scattering is best suited to study isospin violation since the isovector operator $(m_u - m_d)(\bar{u}u - \bar{d}d)$ contributes at the same order as the isoscalar chiral symmetry breaking one, $(m_u + m_d)(\bar{u}u + \bar{d}d)$. In the isospin limit, the πN scattering amplitude decomposes into isoscalar and isovector parts. This allows to formulate a variety of triangle relations between measurable channels that vanish if isospin were an exact symmetry, see e.g.[1]. As a first necessary step, one has to consider the isospin limit. In the framework of heavy baryon CHPT, the S- and P-wave phases can be described precisely in a fourth order calculation [2]. Fixing the low-energy constants (LECs) for pion lab momenta between 50 and 100 MeV by fitting to the existing πN partial wave analyses (KH, VPI/GWU, Zürich), one can predict threshold parameters and the phases at higher energies. Most importantly this calculation shows convergence of the chiral expansion. The inclusion of virtual photons (denoted as γ^*) in the effective πN Lagrangian is straightforward, for a detailed discussion see e.g. [3]. In [4], a complete third order calculation was performed including one- and two-photon exchange diagrams, photon loop corrections, soft photon radiation and three em LECs, which parameterize the effects of hard photons and vacuum polarization. A fit to all data (with the exception of the ones generally considered inconsistent) below pion lab momenta of 100 MeV allowed to determine the strong and em LECs and to uniquely separate the hadronic and the em amplitudes. A very pronounced difference in the hadronic amplitude for elastic scattering $\pi^- p \to \pi^- p$ compared to the standard PWA's was observed. This difference could be traced back to the inclusion of non-linear $\pi\pi\bar{N}N\gamma^*$ couplings as demanded by chiral symmetry. Such effects have to be included and thus a revision of the commonly used em corrections like e.g. the NORDITA approach is called for. In the S–wave triangle relation between elastic scattering, $\pi^\pm p \to \pi^\pm p$, and charge exchange, $\pi^- p \to \pi^0 n$, one finds strong isospin violation of -0.75% in the low-energy region, an order of magnitude smaller than reported in the literature [5] but consistent we the expected size of isospin violation. It is now of utmost importance to map these result obtained in chiral perturbation theory on a dispersive representation to analyze also πN data at higher energies.

[1] N. Fettes, Ulf-G. Meißner and S. Steininger, Phys. Lett. B **451** (1999) 233; N. Fettes and Ulf-G. Meißner, Phys.Rev. C **63** (2001) 045201.

[2] N. Fettes and Ulf-G. Meißner, Nucl. Phys. A **676** (2000) 311.

[3] G. Müller and Ulf-G. Meißner, Nucl. Phys. B **556** (1999) 265.

[4] N. Fettes and Ulf-G. Meißner, Nucl. Phys. A **693** (2001) 693.

[5] W.R. Gibbs, Li Ai, and W.B. Kaufmann, Phys. Rev. Lett. **74** (1995) 3740; E. Matsinos, Phys. Rev. C **56** (1997) 3014.